

Literature review on corneal sensitivity with contact lens wear

Daniela S. Nosch^{1,2}

¹ Ph.D. M.Sc., MCOptom, DipTP(AS), FBCLA FEAAO · ² Institute of Optometry, University of Applied Sciences Northwestern Switzerland, Olten, Switzerland

Received 29 November 2023; accepted 12 February 2024

Abstract

Purpose. To investigate the effect of contact lens (CL) wear on ocular surface sensitivity and to determine and summarise the relationship between corneal sensitivity and CL comfort.

Material and Methods. A literature search was carried out in PubMed, Scopus and Google Scholar, whereby relevant publications were reviewed and summarised during the period of 13th and 20th November 2023.

Results. For the evaluation of the published literature, it is important to acknowledge that corneal sensitivity measurements are influenced by both, the psychophysical technique and the type of instrument used. Study results show a clear but reversible decrease in corneal sensitivity with oxygen-impermeable polymethyl methacrylate (PMMA) contact lens wear, but little or no effect during daily rigid gas permeable (RGP) CL wear. The results for soft CL wear are more complicated: A decrease in corneal sensitivity may be observed during daily wear of hydrogel CLs with low oxygen permeability, however no change or only a small decrease (depending on the applied measurement method) during silicone hydrogel CL wear. Some studies even found a slight sensitisation with this CL material.

Conclusion. Based on the published study results, it is reasonable to assume that hypoxia is the most likely cause of the reduction in corneal sensitivity during daily CL wear. Successful CL wear with materials that have a sufficiently high oxygen transmissibility has a negligible effect on corneal sensitivity. However, an increased sensation of irritation, particularly with symptomatic CL wear may be observed. This could be caused by (subclinical) inflammatory reactions.

Keywords

Corneal sensitivity; contact lens; corneal sensory nerves, aesthesiometry

Introduction

Corneal sensitivity is determined by a neurological response of the superficial nerve fibre endings in the corneal epithelium. They register mechanical, chemical and thermal irritation and thus provide the cornea with an important protective mechanism against harmful influences from the environment. They can be functionally differentiated as follows:^{1,2} Mechanical and electrical stimulation activates the mechanonociceptors, causing a sensation of touch, pain and/or irritation. Polymodal nociceptors react to mechanical, thermal and chemical stimuli, which causes a burning or even stabbing pain. Cold thermoreceptors increase their activity when the temperature on the ocular surface decreases and osmolarity of the tear film increases, which causes a sensation of cooling and dryness. Cold thermoreceptors can be divided into those with a low threshold and high background activity and those with a high threshold and low background activity at normal corneal temperature.^{1,2} Those with low threshold and high background activity are thought to be responsible for the sensation of cooling, while the others cause the sensation of dryness, irritation and/or pain.³ Cold-sensitive thermoreceptors are involved in the regulation of basal tear film production and blinking.^{4,5} The sensory nerves change their activity during inflammatory reactions and tissue damage on the anterior surface of the eye,⁶⁻⁹ This causes sensations of irritation and pain and influences the blink frequency and tear film production rate.^{10,11} An increase in tear film osmolarity stimulates the cold-sensitive and polymodal nociceptors.^{12,13} An inflammatory event results in a sensitisation of polymodal nociceptors, while cold-thermoreceptors are simultaneously inhibited.^{6,7} The nerve endings in the cornea and conjunctiva are connected to the lacrimal glands and the orbicularis oculi muscle via a complex feedback network (activating brainstem control circuits) to monitor and maintain the health of the

anterior surface of the eye and the tear film at all times.¹⁴ They trigger the release of trophic substances (neuropeptides and neurotrophins) to regulate the healing process after injury.¹⁵

Contact lens (CL) wear and the use of care products lead to mechanical forces, temperature fluctuations and chemical stimulation of the ocular surface, either directly through exogenous irritation or indirectly through the release of endogenous agents due to cell damage, hypoxia or changes in pH or osmolarity.¹⁶ These stimuli not only lead to stimulation of the sensory nerves, but also to damage to the nerve endings and local inflammation.¹⁰ These events in turn further activate and sensitise the sensory nerves, leading to discomfort in some CL wearers.¹⁶

Material and methods

A literature search was conducted in PubMed, Scopus and Google Scholar to investigate the influence of CL wear on ocular surface sensitivity. Three searches were conducted using the English keywords "contact lens AND corneal sensitivity", "contact lens AND conjunctival sensitivity" and "contact lens AND lid margin sensitivity" in the period from 13-20 November 2023.

The measurement of corneal sensitivity

The measurement of corneal sensitivity enables the assessment of the functionality of the pain-sensitive superficial corneal nerves. This provides important information about the health of the cornea during the course of a disease pro-




Figure 1: Diagram of the Swiss Liquid Jet Aesthesiometer for Corneal Sensitivity (SLACS)




Figure 2: Example of a liquid jet travelling toward the ocular surface

cess, during the healing phase after an injury or refractive surgery,¹⁷⁻²⁰ as well as during CL wear.¹⁶

With the only commercially available instrument, the Cochet-Bonnet (CB) aesthesiometer, a tactile mechanical stimulus is generated with a 6 cm long nylon thread in 0.12 mm and 0.08 mm thickness: depending on the length of this thread, a more or less intense stimulation is triggered on the corneal surface.²¹ The nylon thread with a thickness of 0.08 mm covers a rather higher sensitivity range, but is unfortunately no longer available. Unfortunately, this instrument has several limitations:²²⁻²⁵ It is an invasive method, which may damage the corneal epithelium; reproducibility is poor because precise centration of the nylon thread is not possible and also the force on the cornea cannot be controlled – even a slight deviation from the correct angle of the thread end to the cornea significantly affects measurement accuracy; the stimulus range is limited, especially in the upper sensitivity range, which means that corneal sensitivity is underestimated and slight sensitivity changes cannot be detected; humidity affects the bending ability of the thread.

To overcome these shortcomings, various prototypes of non-contact air jet aesthesiometers were developed to generate either cooling or warming on the ocular surface, to stimulate temperature-sensitive or mechano- or polymodal nociceptors.²⁶⁻³⁰ However, it is questioned whether the thermal component of the stimulus can be eliminated to produce a true mechanical stimulus, as the air jet causes an evaporative cooling effect on the cornea depending on the stimulus intensity.³¹ It is therefore assumed that the mode of action of this type of stimulus results in both a localised reduction in the superficial ocular surface temperature and a slight indentation of the epithelial surface.^{22,24,32} It is also problematic that the air jet stimulus spreads in a lateral movement over the entire corneal surface, resulting in a stimulus footprint that is difficult to determine.³³ It is unclear to what extent the results of previous studies investigating the CL effect on corneal sensitivity were influenced by device limitations and/or differences in the nature of the stimulus.

Recently, a novel non-invasive liquid jet prototype was developed at the University of Applied Sciences, FHNW

(Switzerland) (Figures 1 and 2) using small droplets of isotonic saline solution with a pH of 7.4 and an osmolarity of 290.2 mOsm/L adapted to the normal tear film: The Swiss Liquid Jet Aesthesiometer for Corneal Sensitivity (SLACS). The liquid jet emerges from a micro-valve equipped with a heating coil and a temperature sensor. The intensity of the stimulus is controlled with variable pressure levels. In contrast to CB and most other aesthesiometer prototypes, this one uses a software algorithm, which means that measurements can be carried out independently of the influence of the examiner. The functional principle and the relevant physical properties of this new prototype were described,³⁴ and it was clinically validated in a study with 90 participants.³⁵ With its wide stimulus range and pressure resolution, SLACS can potentially detect much smaller variations in sensitivity.

The influence of contact lenses on the sensitivity of the cornea and conjunctiva

CL wear naturally interacts with the ocular surface and can affect corneal sensitivity. It is postulated that the following mechanisms cause a decrease in the sensitivity of the ocular surface when wearing CLs: Metabolic impairment of the cornea due to hypoxia (reduced oxygen supply),³⁶⁻⁴⁰ sensory adaptation to mechanical irritation^{41,42} and corneal acidosis.⁴³ Metabolic impairment due to hypoxia may be caused by an impairment in the production of the neurotransmitter acetylcholine, which has a higher concentration in the corneal epithelium than in other areas of the body.⁴⁴ It is therefore assumed that acetylcholine plays an important role in ionic transport (sodium chloride) in the cornea, which in turn has an influence on the generation of nerve impulses.⁴⁵ A sensory adaptation to mechanical stimulation is plausible due to the altered and reversible arrangement of the nerves in the epithelial subbasal nerve plexus during the orthokeratology CL wear.⁴⁶ Small changes in the pH value significantly alter nerve activity,⁴³ a reduction in pH occurs as a result of hypercapnia (accumulation of carbon dioxide). Sensitisation of the corneal nerves, on the other hand, is thought to be the result of hyperosmolarity and/or inflammatory mediators during CL wear.^{27,47} **Table 1** summarises all studies and their results that have investigated the influence of CL wear on the sensitivity of the ocular surface.

The influence of CL wear on corneal sensitivity

Studies using the CB measurement method observed a clear but reversible decrease in corneal sensitivity when wearing oxygen-impermeable polymethyl methacrylate (PMMA) CLs,^{37,48-53} but only a slight effect or no effect at all during daily rigid gas permeable (RGP) CL wear.^{37,38,42,52,54} Even with SLACS, no difference was found between RGP CL wearers and silicone hydrogel (SH) CL wearers and a control group.⁵⁴

Table 1: Summary of published clinical studies on the influence of CL wear on the sensitivity of the ocular surface; SH = silicone hydrogel; CB = Cochet-Bonnet; SLACS = Swiss Liquid Jet Aesthesiometer for Corneal Sensitivity; RGP = rigid gas permeable; PMMA = polymethyl methacrylate

Authors (year of publication)	Sample size; age	Measurement method	Contact lens types	The most important findings
Lowther and Hill (1968) ⁶⁵	n = 4; age 19-21 years	CB	PMMA	Decrease of eyelid margin sensitivity with PMMA KL
Millodot (1974) ³⁶	n = 12; age: 21-27 years	CB (ascending staircase method)	Hydrogel with low oxygen permeability	Lower corneal sensitivity after 8 h of CL wear, compared to baseline
Norn (1975) ⁶²	n = 102; no information on age	CB (ascending staircase method)	PMMA and RGP	lower bulbar conjunctival sensitivity
Millodot (1976) ⁴⁸	n = 12, age: 21-31 years; 7 women	CB (ascending staircase method)	PMMA	Gradually lower corneal sensitivity after 4, 8 and 12 hours of CL wear, compared to baseline; high variability
Millodot (1977, 1978) ^{49,50}	n = 82 and n = 91, age: 21-46 years	CB (ascending staircase method)	PMMA	Decrease in corneal sensitivity dependent on duration of CL wear in years and recovery after cessation of CL wear
Millodot (1979) ⁵¹	n = 9	CB (ascending staircase method)	PMMA and CAB	Significant corneal sensitivity loss with PMMA and slightly less pronounced with CAB CLs after 10 h of wear, compared to baseline
Douthwaite and Connelly (1986) ⁵²	n = 76; age: 15-59 years; 56 women;	CB (ascending staircase method)	PMMA and RGP	Significantly more pronounced loss of corneal sensitivity after 3 months of PMMA than with RGP CL wear
Bergenske and Polse (1987) ³⁷	n = 10; average age: 40.1; 27-55 years; 9 women	CB (ascending staircase method)	PMMA → newly fitted with RGP	Loss of corneal sensitivity with PMMA CLs, which recovers after switching to RGP CLs
Sanaty and Temel (1998) ⁵³	n = 20; average age: 29.7; 21-45 years	CB (ascending staircase method)	PMMA	Reversible corneal sensitivity loss with PMMA CL wear
Larke and Hirji (1979) ⁵⁵	n = 57	CB (ascending staircase method)	Hydrogel with low oxygen permeability	Gradual loss of corneal sensitivity in newly fitted hydrogel CLs over the course of 20 weeks
Velasco et al. (1994) ³⁹	n = 40 in group 1 and n = 27 in group 2 of CL wearers	CB (ascending staircase method)	Hydrogel: Group 1 with 38%, Group 2 with 55% water content	Lower corneal sensitivity in both CL groups after 8 h wearing time, compared to baseline; more pronounced in group 1 with low water content and oxygen permeability
Murphy et al. (2001) ³⁸	n = 40 RGP, n = 40 hydrogel and n = 40 control group; age: 34; 19-68 years (total CL group); 50 women	Non-contact corneal aesthesiometer (NCCA): air jet method (double staircase method)	RGP and hydrogel	lower corneal sensitivity with RGP and hydrogel CL wear than in the control group; no difference between the CL types

Table 1 (Continued)

Authors (year of publication)	Sample size; age	Measurement method	Contact lens types	The most important findings
Stapleton et al. (2004) ²⁷	n = 10; average age: 24 years; 21-30 years; 5 women	CRCERT-Belmonte aesthesiometer; air jet (34 °C; staircase method)	Hydrogel and SH	No change in corneal sensitivity and no difference between hydrogel and SH; increased bulbar conjunctival sensitivity with SH CLs
Hiraoka et al. (2009) ⁶¹	n = 17; average age: 23.5 ± 3.2 years	CB (ascending staircase method)	RGP (Orthokeratology)	lower corneal sensitivity three months after fitting Orthokeratology CLs
Situ et al. (2010) ⁴⁷	n = 50; average age: 25.2; 18-45 years; 35 women	CB (ascending staircase method) and Belmonte aesthesiometer with air jet (50 °C)	SH	Lower corneal sensitivity with CB CLs and air jet aesthesiometry and higher bulbar conjunctival sensitivity with air jet aesthesiometer, but not with CB
Golebiowski et al. (2012) ⁵⁶	n = 27; average age: 40 years; 28-52 years	CRCERT-Belmonte aesthesiometer; air jet (34 °C; staircase method)	Hydrogel with deep oxygen permeability → newly supplied with SH	No difference in corneal sensitivity between overnight hydrogel CL wear and control group, but lower corneal sensitivity after cessation of hydrogel CL wear, which did not change with subsequent SH CL wear; no difference/change in relation to the bulbar conjunctiva
Lum et al. (2013) ⁴²	n = 20; average age: 24; 19-39 years	CB (ascending staircase method) and NCCA (double staircase method)	SH, RGP and Orthokeratology CLs successively fitted on all participants	Lower corneal sensitivity after orthokeratology, no change with RGP and SH CLs
Hiscox et al. (2015) ⁶⁶	n = 15 in CL group, n = 15 in control group;	CB (ascending staircase method)	Soft CLs (material not specified)	No difference in eyelid margin sensitivity compared to the control group
Navascues-Cornago et al. (2015) ⁶⁴	n = 35 in CL group (25.6 ± 7.4 years), n = 35 in control group (27.7 ± 7.3 years)	CB (ascending staircase method)	Soft CLs (material not specified)	Lower sensitivity at the inferior bulbar conjunctiva in the CL group after 12 h of CL wear and compared to baseline; higher lid margin sensitivity in the CL group after 12 h of CL wear compared to the control group
Igarashi et al. (2015) ⁶³	n = 12 in soft CL, n = 14 in control group; average age: 28.3 ± 4.6 years	CB (ascending staircase method)	Soft CLs (material not specified)	Increase in sensitivity of the bulbar and tarsal conjunctiva in the CL group
Nosch et al. (2019) ⁶⁸	n = 34; average age 23.85 ± 5.39 years; 17 women	CB (double staircase method)	RGP	Lower lid margin sensitivity after short term RGP CL wear for a duration of 45 min
Nosch et al. (2023) ⁵⁴	n = 33 in SH group (age: 27.42 ± 6.83 years; 17 women); n = 30 in RGP group (36.90 ± 9.68 years; 21 women); n = 33 in control group (26.06 ± 6.19 years; 23 women)	CB and SLACS (each with double staircase method)	SH, RGP and control group	No statistically significant difference in sensitivity between the three groups with either measurement methods

Table 1 (Continued)

Authors (year of publication)	Sample size; age	Measurement method	Contact lens types	The most important findings
Kleinschmidt and Zimmermann (2022) ⁵⁸	n = 30; average age: 26.93 ± 9.82 years; 22 women	CB and SLACS (each with double staircase method)	SH	Increased corneal sensitivity with SLACS, but not with CB, compared to baseline
Seghetti (2023) ⁵⁷	n = 38; average age: 26.55 ± 5.7 years; 26 women	SLACS (double staircase method)	SH	No statistically significant difference in sensitivity
Angel et al. (2023) ⁷⁷	n = 14 in symptomatic CL group; n = 17 in asymptomatic CL group; n = 29 in control group (age: 24.5 ± 0.8 years); total age in CL group: 23.8 ± 1 years	CRCERT-Belmonte aesthesiometer with air jet stimulus (34 °C): mechanical, chemical and cooling; suprathreshold assessment	Soft CLs (hydrogel and SH)	Higher sensation of irritation in the CL group than in the control group, especially in asymptomatic CL group; lower perception of stimulus intensity in the symptomatic CL group

The results to date for soft CL wear are more complicated: the studies conducted with CB agree that corneal sensitivity decreases with hydrogel CLs with a low Dk value (= low oxygen permeability),^{36,39,55} but not or only to a small extent with SH-CLs with a high Dk value (= high oxygen permeability).^{42,47} Studies with air jet aesthesiometry and SLACS observed no or only very minor effects in hydrogel and SH-CL wearers with a low or high Dk value.^{27,42,47,54,56,57} Interestingly, however, in a study with SLACS (but not with CB), a slight sensitisation of the cornea was observed in SH-CL with overnight wear after one week compared to baseline.⁵⁸ This increased nerve activity could be an expression of a subclinical inflammatory reaction or indicate a certain biochemical stress.

Previously published studies and a literature review by Stapleton et al.⁵⁹ support the hypothesis that hypoxia is primarily responsible for a decrease in corneal sensitivity when wearing oxygen-impermeable PMMA CLs rather than sensory adaptation to mechanical stimuli: Corneal sensitivity recovered when switching from PMMA to RGP CLs.³⁷ When wearing RGP CLs, only a very slight⁵² or no decrease in sensitivity^{42,54} was observed compared to a control group^{37,42} or a soft CL group.

A hypoxic aetiology is confirmed by the decrease in sensitivity overnight without CL wear^{60,61} and the reduced corneal sensitivity a closed eyelid.⁴⁴

With regard to daily RGP CL wear, the results are promising, as a reduction in corneal sensitivity due to a delayed and/or reduced response of the superficial nerve endings could potentially lead to an increased risk of infection. The short-term and reversible reduction in corneal sensitivity with orthokeratology is likely to be due to the mechanically altered arrangement of nerve fibres in the epithelial subbasal nerve plexus rather than hypoxia.^{42,46,62}

The influence of CL wear on the bulbar conjunctiva

Only few studies looked at the influence of CL wear on the sensitivity of the bulbar conjunctiva and it is therefore unclear whether changes in sensitivity are to be expected: A study conducted with CB observed lower sensitivity with PMMA and RGP CL wear.⁶³ Using air-jet aesthesiometry, two studies found a sensitisation of the bulbar conjunctiva when wearing SH-CLs,^{27,47} using CB, one study found no change when wearing soft CLs,⁴⁷ another, however, observed sensitisation.⁶⁴ In contrast, a lower sensitivity of the inferior bulbar conjunctiva was noted after 12 months of wearing soft CLs compared to a control group and to baseline.⁶⁵ In contrast, another study found no difference in long-term overnight hydrogel CL wearers with air-jet aesthesiometry compared to a control group.⁵⁶

The influence of CL wear on the tarsal conjunctiva

According to a study from the 1960s with a very small sample of four participants, the sensitivity of the lid margin (measured with CB) decreases when wearing PMMA, RGP and hydrogel CLs with a low Dk value.⁶⁶ In contrast, a more recent study observed no change in lid margin sensitivity when wearing modern soft lenses (material not specified),⁶⁷ another reported a higher lid margin sensitivity compared to a control group.⁶⁴ Another study also found a sensitisation in the CL group after 12 h of wearing modern soft CLs (materials not specified) compared to the control group.⁶⁵ This could be caused by a lack of wearing comfort due to interaction between the lid margin and the CL surface. In contrast, another study confirmed a lower lid margin sensitivity after wearing

RGP CLs once for 45 min.⁶⁸ In the search for an answer to the question of which individuals are likely to have fewer adaptation problems with RGP CLs, it also investigated whether spontaneous wearing comfort could correlate with eyelid margin sensitivity at baseline or with the decrease in sensitivity during RGP CL wear. Unfortunately, no correlation could be established. Sensitivity measurements on the eyelid are challenging because they require ectropionisation, which is an unnatural situation for the eye.

When assessing the results of all these studies, it must be borne in mind that the measurement of corneal sensitivity is influenced by both the psychophysical technique and the type of instrument used. Unfortunately, as already mentioned, the only commercially available instrument, the tactile CB aesthesiometer, is unsuitable for everyday clinical use.

Various prototypes have been developed with air jet stimuli to generate either cooling or warming of the ocular surface that stimulate temperature- or mechanically-sensitive or polymodal nociceptors.²⁶⁻³⁰ However, it is controversial whether the thermal component of the stimulus can be eliminated to produce a true mechanical stimulus, as the air jet produces an evaporative cooling effect on the moist cornea depending on the airflow intensity.³¹ It is therefore assumed that the mode of action of this type of stimulus causes both local cooling and indentation of the epithelial surface.^{22,24,32} Another problem is that the air stimulus spreads in a lateral movement over the entire ocular surface, resulting in a stimulus footprint that is difficult to determine.³³ The SLACS prototype was developed to overcome this challenge using a small liquid jet that can be adjusted to the temperature of the ocular surface. Unlike CB and most other aesthesiometer prototypes, it uses a software algorithm that is independent of the examiner.

The influence of CL care products

According to studies, care products preserved with polyhexanide in combination with certain SH-CL materials can cause corneal and conjunctival staining as well as bulbar hyperaemia and CL discomfort.⁶⁹⁻⁷² In a crossover pilot study with the CB aesthesiometer, Epstein found reduced corneal sensitivity with polyhexanide-preserved care products compared to polyquad-preserved care products.⁷³ However, these differences were not statistically significant, possibly due to the small sample size. Situ et al. also investigated the effects of different SH CL/care product combinations on the sensitivity of the ocular surface in a larger group of 48 subjects using CB and the air jet anaesthesiometer:⁴⁷ They compared the sensitivity changes in CL wearers using hydrogen peroxide-based care products and two different multipurpose solutions (preserved with polyhexanide and polyquad/Aldox). They obtained reduced corneal and conjunctival thresholds for chemical sensitivity with the polyhexanide-preserved care product compared to the one preserved with polyquad/Aldox. However, this difference was only statistically significant for the corneal measurements. No statistical differences were found with regard to the tactile and pneumatic mechanical threshold values. They attributed the increased chemical

sensitivity using the polyhexanide-preserved care product to the increased prevalence of corneal and conjunctival staining, which could have resulted in activation of the polymodal nociceptors.

What role do the superficial corneal nerves play in CL comfort?

When wearing CLs, a complex and multifactorial stimulation of the functionally different nerve endings is triggered. Impaired comfort is caused by mechanical, altered osmolarity, cooling and/or chemical effects.¹⁶

Due to their direct interaction with the front surface of the eye, CLs may cause mechanical irritation. Friction with the ocular is caused by the edge properties of the CLs (rounded or sharp-edged), the rigidity and the surface properties. The tear film, already thinned by the presence of the CL, continues to decrease in thickness, blink frequency increases, resulting in eyelid wiper epitheliopathy⁷⁴ and, due to the increased shear forces during a blink, lid-parallel conjunctival folds may also form.⁷⁵ In this situation, the polymodal nociceptors and the mechanoreceptors are stimulated, which leads to a sensation of irritation and foreign body sensation. Inadequate sensory adaptation to CLS also causes discomfort.⁷⁶ Another consequence of these processes is hyperosmolarity, which stimulates the polymodal and cold-sensitive nociceptors, causing a sensation of dryness, burning and cooling. In this context, inflammatory mediators are also released, which in turn sensitise polymodal nerve endings in the cornea and conjunctiva, which also results in irritation and burning.

The difference between symptomatic and asymptomatic CL wearers has been little researched to date: One study showed that symptomatic CL wearers reacted more sensitively to suprathreshold stimuli.⁷⁶ According to another study, corneal sensitivity decreases with increasing symptoms.⁷⁷ A recently published study compared the subjective perception of intensity and irritation to suprathreshold mechanical, chemical and cooling stimuli in an asymptomatic, symptomatic CL- and control group using an air jet aesthesiometer.⁷⁸ They observed a higher sensation of irritation in the CL- than in the control group, especially in asymptomatic CL wearers. In contrast, they recorded a lower intensity sensation to cooling irritation in the symptomatic CL group. They postulate that sensitisation of nociceptors causes irritation on an inflamed and poorly wetting cornea. At the same time, the cold-sensitive nociceptors would reduce their activity due to the inflammatory reaction. In view of the rather small sample with different group sizes, however, these results must be confirmed in a larger-scale study.

In summary, based on studies published to date, hypoxia is the most likely cause of a mechanical reduction in corneal sensitivity during daytime CL wear. Successful daily wear of modern, sufficiently oxygen-permeable CLs has a negligible effect on corneal sensitivity. However, there may be an increased sensation of irritation, particularly with symptomatic CL wear. It is reasonable to assume that (subclinical)

inflammatory reactions play a significant role. Further studies are needed to investigate the role of the bulbar conjunctiva and the lid margin (especially in the region of the lid wiper) in comfort problems with modern CL materials, as this is where most interaction with the CL occurs.

Author



Prof. Dr. Daniela Nosch

E-Mail:
daniela.nosch@fhnw.ch

References

- Gallar, J., Pozo, M. A., Tuckett, R. P., Belmonte, C. (1993). Response of sensory units with unmyelinated fibres to mechanical, thermal and chemical stimulation of the cat's cornea. *J. Physiol.*, 468, 609–622.
- González-González, O., Bech, F., Gallar, J., Merayo-Llodes, J., Belmonte, C. (2017). Functional properties of sensory nerve terminals of the mouse cornea. *Invest. Ophthalmol. Vis. Sci.*, 58, 404–415.
- Acosta, M. C., Belmonte, C., Gallar, J. (2001). Sensory experiences in humans and single-unit activity in cats evoked by polymodal stimulation of the cornea. *J. Physiol.*, 534, 511–525.
- Parra, A., Madrid, R., Echevarria, D., del Olmo, S., Morenilla-Palao, C., Acosta, M. C., Gallar, J., Dhaka, A., Viana, F., Belmonte, C. (2010). Ocular surface wetness is regulated by TRPM8-dependent cold thermoreceptors of the cornea. *Nat. Med.*, 16, 1396–1399.
- Quallo, T., Vastani, N., Horridge, E., Gentry, C., Parra, A., Moss, S., Viana, F., Belmonte, C., Andersson, D. A., Bevan, S. (2015). TRPM8 is a neuronal osmosensor that regulates eye blinking in mice. *Nat. Commun.*, 6, 7150.
- Acosta, M. C., Luna, C., Quirce, S., Belmonte, C., Gallar, J. (2013). Changes in sensory activity of ocular surface sensory nerves during allergic keratoconjunctivitis. *Pain*, 154, 2353–2362.
- Acosta, M. C., Luna, C., Quirce, S., Belmonte, C., Gallar, J. (2014). Corneal sensory nerve activity in an experimental model of UV keratitis. *Invest. Ophthalmol. Vis. Sci.*, 55, 3403–3412.
- Kovács, I., Luna, C., Quirce, S., Mizerska, K., Callejo, G., Riestra, A., Fernández-Sánchez, L., Meseguer, V. M., Cuenca, N., Merayo-Llodes, J., Acosta, M. C., Gasull, X., Belmonte, C., Gallar, J. (2016). Abnormal activity of corneal cold thermoreceptors underlies the unpleasant sensations in dry eye disease. *Pain*, 157, 399–417.
- Luna, C., Mizerska, K., Quirce, S., Belmonte, C., Gallar, J., Acosta, M. D. C., Meseguer, V. (2021). Sodium channel blockers modulate abnormal activity of regenerating nociceptive corneal nerves after surgical lesion. *Invest. Ophthalmol. Vis. Sci.*, 62, 2.
- Belmonte, C., Acosta, M. C., Merayo-Llodes, J., Gallar, J. (2015). What causes eye pain? *Curr. Ophthalmol. Rep.*, 3, 111–121.
- Belmonte, C. (2019). Pain, dryness, and itch sensations in eye surface disorders are defined by a balance between inflammation and sensory nerve injury. *Cornea*, 38, Suppl. 1, S11–S24.
- Parra, A., Gonzalez-Gonzalez, O., Gallar, J., Belmonte, C. (2014). Tear fluid hyperosmolality increases nerve impulse activity of cold thermoreceptor endings of the cornea. *Pain*, 155, 1481–1491.
- Belmonte, C., Nichols, J. J., Cox, S. M., Brock, J. A., Begley, C. G., Bereiter, D. A., Dartt, D. A., Galor, A., Hamrah, P., Ivanusic, J. J., Jacobs, D. S., McNamara, N. A., Rosenblatt, M. I., Stapleton, F., Wolffsohn, J. S. (2017). TFOS DEWS II pain and sensation report. *Ocul. Surf.*, 15, 404–437.
- Stern, M. E., Gao, J., Siemasko, K. F., Beuerman, R. W., Pflugfelder, S. C. (2004). The role of the lacrimal functional unit in the pathophysiology of dry eye. *Exp. Eye Res.*, 78, 409–416.
- Belmonte, C., Garcia-Hirschfeld, J., Gallar, J. (1997). Neurobiology of ocular pain. *Prog. Retin. Eye Res.*, 16, 117–156.
- Stapleton, F., Marfurt, C., Golebiowski, B., Rosenblatt, M., Bereiter, D., Begley, C., Dartt, D., Gallar, J., Belmonte, C., Hamrah, P., Willcox, M. (2013). The TFOS International Workshop on Contact Lens Discomfort: Report of the subcommittee on neurobiology. *Invest. Ophthalmol. Vis. Sci.*, 54, TFOS71–97.
- Gallar, J., Acosta, M. C., Moilanen, J. A. O., Holopainen, J. M., Belmonte, C., Tervo, T. M. T. (2004). Recovery of corneal sensitivity to mechanical and chemical stimulation after laser in situ keratomileusis. *J. Refract. Surg.*, 20, 229–235.
- Pérez-Santonja, J. J., Sakla, H. F., Cardona, C., Chipont, E., Alió, J. L. (1999). Corneal sensitivity after photorefractive keratectomy and laser in situ keratomileusis for low myopia. *Am. J. Ophthalmol.*, 127, 497–504.
- Murphy, P. J., Corbett, M. C., O'Brart, D., Verma, S., Patel, S., Marshall, J. (1999). Loss and recovery of corneal sensitivity following photorefractive keratectomy for myopia. *J. Refract. Surg.*, 15, 38–45.
- Darwish, T., Brahma, A., O'Donnell, C., Efron, N. (2007). Subbasal nerve fiber regeneration after LASIK and LASEK assessed by noncontact esthesiometry and in vivo confocal microscopy: Prospective study. *J. Cataract Refract. Surg.*, 33, 1515–1521.
- Cochet, P., Bonnet, R. (1961). L'esthésiométrie cornéenne. Réalisation et intérêt pratique. *Bull. Ophthalmol. Fr.*, 6, 541–550.
- Murphy, P. J., Lawrenson, J. G., Patel, S., Marshall, J. (1998). Reliability of the Non-Contact Corneal Aesthesiometer and its comparison with the Cochet-Bonnet aesthesiometer. *Ophthalmic Physiol. Opt.*, 18, 532–539.
- Millodot, M., Larson, W. (1967). Effect of bending of the nylon thread of the Cochet-Bonnet aesthesiometer upon the recorded pressure. *Die Kontaktlinse*, 1, 5–6.
- Golebiowski, B., Papas, E., Stapleton, F. (2011). Assessing the sensory function of the ocular surface: Implications of use of a non-contact air jet aesthesiometer versus the Cochet-Bonnet aesthesiometer. *Exp. Eye Res.*, 92, 408–413.
- Lum, E., Murphy, P. J. (2018). Effects of ambient humidity on the Cochet-Bonnet aesthesiometer. *Eye (Lond)*, 32, 1644–1651.
- Belmonte, C., Acosta, M., Schmelz, M., Gallar, J. (1999). Measurement of corneal sensitivity to mechanical and chemical stimulation with a CO₂ aesthesiometer. *Invest. Ophthalmol. Vis. Sci.*, 40, 513–519.
- Stapleton, F. (2004). Corneal and conjunctival sensitivity to air stimuli. *Brit. J. Ophthalmol.*, 88, 1547–1551.
- Feng, Y., Simpson, T. L. (2003). Nociceptive sensation and sensitivity evoked from human cornea and conjunctiva stimulated by CO₂. *Invest. Ophthalmol. Vis. Sci.*, 44, 529–532.
- Murphy, P., Morgan, P., Patel, S., Marshall, J. (1996). A new non-contact corneal aesthesiometer. *Ophthalmic Physiol. Opt.*, 16, 101–107.
- Vega, J. A., Simpson, T. L., Fonn, D. (1999). A noncontact pneumatic esthesiometer for measurement of ocular sensitivity: A preliminary report. *Cornea*, 18, 675–681.
- Nosch, D. S., Pult, H., Albon, J., Purslow, C., Murphy, P. J. (2018). Does air gas aesthesiometry generate a true mechanical stimulus for corneal sensitivity measurement? *Clin. Exp. Optom.*, 101, 193–199.
- Murphy, P. J., Morgan, P. B., Patel, S., Marshall, J. (1999). Corneal surface temperature change as the mode of stimulation of the Non-Contact Corneal Aesthesiometer. *Cornea*, 18, 333–342.
- Golebiowski, B., Lim, M., Papas, E., Stapleton, F. (2013). Understanding the stimulus of an air-jet aesthesiometer: computerised modelling and subjective interpretation. *Ophthalmic Physiol. Opt.*, 33, 104–113.
- Nosch, D. S., Oscity, M., Steigmeier, P., Käser, E., Loepfe, M., Joos, R. E. (2022). Working principle and relevant physical properties of the Swiss Liquid Jet Aesthesiometer for Corneal Sensitivity (SLACS) evaluation. *Ophthalmic Physiol. Opt.*, 42, 609–618.
- Nosch, D. S., Käser, E., Bracher, T., Joos, R. E. (2023). Clinical application of the new Swiss Liquid Jet Aesthesiometer for corneal sensitivity measurement. *Clin. Exp. Optom.*, 107, 14–22.
- Millodot, M. (1974). Effect of soft lenses on corneal sensitivity. *Acta Ophthalmol. (Copenh)*, 52, 603–608.
- Bergenske, P. D., Polse, K. A. (1987). The effect of rigid gas permeable lenses on corneal sensitivity. *J. Am. Optom. Assoc.*, 58, 212–215.
- Murphy, P. J., Patel, S., Marshall, J. (2001). The effect of long-term, daily contact lens wear on corneal sensitivity. *Cornea*, 20, 264–269.
- Velasco, M. J., Bermúdez, F. J., Romero, J., Hita, E. (1994). Variations in corneal sensitivity with hydrogel contact lenses. *Acta Ophthalmol. (Copenh)*, 72, 53–56.
- Millodot, M., O'Leary, D. J. (1980). Effect of oxygen deprivation on corneal sensitivity. *Acta Ophthalmol. (Copenh)*, 58, 434–439.

- 41 Polse, K. A. (1978): Etiology of corneal sensitivity changes accompanying contact lens wear. *Invest. Ophthalmol. Vis. Sci.*, 17, 1202–1206.
- 42 Lum, E., Golebiowski, B., Gunn, R., Babhoota, M., Swarbrick, H. (2013). Corneal sensitivity with contact lenses of different mechanical properties: *Optom. Vis. Sci.*, 90, 954–960.
- 43 Brennan, N. A., Bruce, A. S. (1991). Esthesiometry as an indicator of corneal health. *Optom. Vis. Sci.*, 6, 699–702.
- 44 Millodot, M., O’Leary, D. J. (1979). Loss of corneal sensitivity with lid closure in humans. *Exp. Eye Res.*, 29, 417–421.
- 45 Pesin, S. R., Candia, O. A. (1982). Acetylcholine concentration and its role in ionic transport by the corneal epithelium. *Invest. Ophthalmol. Vis. Sci.*, 22, 651–659.
- 46 Lum, E., Golebiowski, B., Swarbrick, H. A. (2012). Mapping the corneal sub-basal nerve plexus in orthokeratology lens wear using in vivo laser scanning confocal microscopy. *Invest. Ophthalmol. Vis. Sci.*, 53, 1803–1809.
- 47 Situ, P., Simpson, T. L., Jones, L. W., Fonn, D. (2010). Effects of silicone hydrogel contact lens wear on ocular surface sensitivity to tactile, pneumatic mechanical, and chemical stimulation. *Invest. Ophthalmol. Vis. Sci.*, 51, 6111–6117.
- 48 Millodot, M. (1976). Effect of length of wear of contact lenses on corneal sensitivity. *Acta Ophthalmol. (Copenh)*, 54, 721–730.
- 49 Millodot, M. (1977). Does the long term wear of contact lenses produce a loss of corneal sensitivity? *Experientia*, 33, 1475–476.
- 50 Millodot, M. (1978). Effect of long-term wear of hard contact lenses on corneal sensitivity. *Arch. Ophthalmol.*, 96, 1225–1227.
- 51 Millodot, M., Henson, D. B., O’Leary, D. J. (1979). Measurement of corneal sensitivity and thickness with PMMA and gas-permeable contact lenses. *Optom. Vis. Sci.*, 56, 628–632.
- 52 Douthwaite, W. A., Connelly, A. T. (1986). The effect of hard and gas permeable contact lenses on refractive error, corneal curvature, thickness and sensitivity. *J. Br. Contact. Lens Assoc.*, 9, 14–20.
- 53 Sanaty, M., Temel, A. (1998). Corneal sensitivity changes in long-term wearing of hard polymethylmethacrylate contact lenses. *Ophthalmologica*, 212, 328–330.
- 54 Nosch, D. S., Käser, E., Christen, A., Schinzel, J., Joos, R. E. (2023). Corneal sensitivity in silicone hydrogel and rigid gas permeable contact lens wear. *Cont. Lens Anterior Eye*, 46, 101888.
- 55 Larke, J. R., Hirji, N.K. (1979). Some clinically observed phenomena in extended contact lens wear. *Br. J. Ophthalmol.*, 63, 475–477.
- 56 Golebiowski, B., Papas, E. B., Stapleton, F. (2012). Corneal and conjunctival sensory function: The impact on ocular surface sensitivity of change from low to high oxygen transmissibility contact lenses. *Invest. Ophthalmol. Vis. Sci.*, 53, 1177–1181.
- 57 Seghetti, M. (2023). Corneal sensitivity in new silicone hydrogel contact lens wearers. October, MSc - Thesis, School of Optometry and Vision Sciences, Cardiff University, Cardiff, UK.
- 58 Kleinschmidt, V., Zimmermann, J. (2023). Changes in corneal sensitivity in extended wear of silicone hydrogel contact lenses. August, BSc-Thesis, Institut für Optometrie, FHNW, Olten, Switzerland.
- 59 Stapleton, F., Chao, C., Golebiowski, B. (2019). Topical review: effects of contact lens wear on corneal, conjunctival, and lid margin sensitivity. *Optom. Vis. Sci.*, 96, 790–801.
- 60 du Toit, R., Vega, J. A., Fonn, D., Simpson, T. (2003). Diurnal Variation of Corneal Sensitivity and Thickness. *Cornea*, 22, 205–209.
- 61 Millodot, M. (1972). Diurnal variation of corneal sensitivity. *Br. J. Ophthalmol.*, 56, 844–847.
- 62 Hiraoka, T., Kaji, Y., Okamoto, F., Oshika, T. (2009). Corneal sensation after overnight orthokeratology. *Cornea*, 28, 891–895.
- 63 Norn, M. S. (1975). Conjunctival sensitivity in pathological cases, with simultaneous measurement of corneal and lid margin sensitivity. *Acta Ophthalmol. (Copenh)*, 53, 450–457.
- 64 Igarashi, T., Ono, M., Fujimoto, C., Suzuki, H., Takahashi, H. (2015). The conjunctival sensitivity in soft contact lens wearers. *Int. Ophthalmol.*, 35, 569–573.
- 65 Navascues-Cornago, M., Morgan, P. B., Maldonado-Codina, C. (2015). Lid margin sensitivity and staining in contact lens wear versus no lens wear. *Cornea*, 34, 808–816.
- 66 Lowther, G. E., Hill, R. M. (1968). Sensitivity threshold of the lower lid margin in the course of adaptation to contact lenses. *Optom. Vis. Sci.*, 45, 587–594.
- 67 Hiscox, R., Evans, K., North, R., Allen, L., Purslow, C. (2015). Examining the effect of soft contact lens wear on lid margin sensitivity. *Cont. Lens Anterior Eye*, 38, e36.
- 68 Nosch, D. S., Joos, R. E., Müller, D., Matt, S. M. (2019). General pain perception sensitivity, lid margin sensitivity and gas permeable contact lens comfort. *Clin. Exp. Optom.*, 103, 766–771.
- 69 Andrasko, G., Ryen, K. (2008). Corneal staining and comfort observed with traditional and silicone hydrogel lenses and multipurpose solution combinations. *Optometry*, 79, 444–454.
- 70 Santodomingo-Rubido, J., Barrado-Navascués, E., Rubido-Crespo, M-J., Sugimoto, K., Sawano, T. (2008). Compatibility of two new silicone hydrogel contact lenses with three soft contact lens multipurpose solutions. *Ophthalmic Physiol Opt.*, 28, 373–381.
- 71 Papas, E. B., Carnt, N., Willcox, M. D. P., Holden, B. A. (2007). Complications associated with care product use during silicone daily wear of hydrogel contact lens: *Eye Contact Lens*, 33, 392–393.
- 72 Jones, L., Macdougall, N., Sorbara, L. G. (2002). Asymptomatic corneal staining associated with the use of balafilcon silicone-hydrogel contact lenses disinfected with a polyaminopropyl biguanide-preserved care regimen: *Optom. Vis. Sc.*, 79, 753–761.
- 73 Epstein, A. B. (2006). Contact lens care products effect on corneal sensitivity and patient comfort: *Eye Contact Lens*, 32, 128–132.
- 74 Shiraishi, A., Yamaguchi, M., Ohashi, Y. (2014). Prevalence of upper- and lower-lid-wiper epitheliopathy in contact lens wearers and non-wearers: *Eye Contact Lens*, 40, 220–224.
- 75 Pult, H., Purslow, C., Murphy, P. J. (2011). The relationship between clinical signs and dry eye symptoms. *Eye (Lond)*, 25, 502–510.
- 76 Chen, J., Simpson, T. L. (2011). A role of corneal mechanical adaptation in contact lens-related dry eye symptoms. *Invest. Ophthalmol. Vis. Sci.*, 52, 1200–1205.
- 77 Martín-Montañez, V. (2015). End-of-day dryness, corneal sensitivity and blink rate in contact lens wearers. *Cont. Lens Anterior Eye*, 38, 148–151.
- 78 Pastor-Zaplana, J. Á., Gallar, J., Acosta, M. C. (2023). Functional changes of the ocular surface sensory nerves due to contact lens use in young symptomatic and asymptomatic users. *Invest. Ophthalmol. Vis. Sci.*, 64, 12.